



2

DTIC
ELECTE
JUN 03 1986
S D D

LABORATORY NOTE NO. 86-57

COMPUTER VISUAL SIMULATION OF CONTRAST SENSITIVITY DEFICITS
INDUCED BY LASER AND CHEMICAL ANTIDOTE EXPOSURE

AD-A168 608

HARRY ZWICK, PhD
DAN MONROE, CPT, MSC
and
LARRY SHERMAN, BS

DTIC
ELECTE
JUN 05 1986
S D D

DIVISION OF OCULAR HAZARDS
and
INFORMATION SCIENCES GROUP

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

DTIC FILE COPY

MARCH 1986

LETTERMAN ARMY INSTITUTE OF RESEARCH
PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

00 0

4 050

4

Computer Visual Simulation of Contrast Sensitivity Deficits Induced by
Laser and Chemical Antidote Exposure--Zwick, Monroe, and Sherman

Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

Edum S. Seatrie 7 May 66
(Signature and date)

This document has been approved for public release and sale; its distribution is unlimited.

Computer Visual Simulation of Contrast Sensitivity Deficits
Induced by Laser and Chemical Antidote Exposure

H.Zwick, PHD, DAC, D.Monroe, CPT, MSC, L.Sherman, DAC
Letterman Army Institute of Research
Presidio of San Francisco, Ca 94129

SUMMARY

Effective training of a complex combat-related task presently requires training that may produce a transient degree of visual impairment. The present paper presents an alternative method, i.e., simulating visual impairment produced by potential combat conditions. The degree of realism provided by such simulation offers development of safer as well as more realistic training techniques.

The ability to represent a complex visual image along a unitary dimension and relate this dimension to target acquisition criteria is critical to military training concepts. Recent experiments have suggested that the physics of form and human spatial vision are uniquely related to the spatial frequency domain (1). Small changes in the amplitude of selective spatial frequencies can have significant effects on human form vision (2).

A scheme to represent a complex target along a single, spatial frequency continuum was first proposed by Johnson (3). The Johnson criterion relates the number of cycles in a square wave grating (black bars periodically alternating with white bars) required for either detection, identification, or recognition, to complex military targets as a function of target range. Complex military targets can be scaled with respect to the number of cycles required for either detection, identification, or recognition.

While the Johnson criterion is adequate for scaling high contrast stimuli in a reliable manner, it is neither parsimonious with current physiological mechanisms underlying human spatial vision nor adaptable to characterizing how the perception of complex images might be degraded by exposure to noxious battlefield conditions.

Many recent experiments reporting such effects indicate that mechanisms underlying spatial vision are selectively altered with regard to their ability to represent the spatial frequency domain. Such degradation in the neural mechanism of spatial vision is not easily represented by changes in the number of cycles required for a visual response.



Availability Codes	
Dist	Avail and/or Special
A-1	

Image degradation that might be produced by potential battlefield exposure conditions can be simulated by modulating the spatial frequency content of a complex image in accordance with the actual amplitude modulation induced by experimental procedures. In this paper, spatial frequency contributions have been modified according to the results of experimental data, to obtain the inverse fourier transform of a complex mage.

METHODS

A computer interfaced with a video frame buffer was used to produce fourier filtered images. The frame buffer used stored images in a 256 by 256 pixel format with 64 shades of grey (4). A two-dimensional fast fourier transform (FFT) was used to alter the spatial frequency content of a complex image in accordance with changes that had been induced in contrast sensitivity experiments. The original image was filtered by reducing the signal amplitude contributions at selected spatial frequencies in direct proportion to losses obtained in contrast sensitivity at corresponding spatial frequency points. An inverse FFT was then performed and the filtered image displayed on a CRT and stored on hard disk.

Data from three contrast sensitivity experiments were used. The acute effects of Q-switched pulsed laser exposure(5), benactazine (6), and atropine (7) on contrast sensitivity were used to obtain degraded images in correspondence with previously obtained contrast sensitivity data.

RESULTS

Contrast sensitivity data, replotted as a percentage of the baseline contrast sensitivity for acute laser exposure (5) and for benactazine (6) are shown in Figure 1. Changes induced by atropine (7) were very slight, varying between 9% and 98% of baseline contrast sensitivity measured over the same spatial frequency range as used for benactazine.




FIGURE 1. Contrast sensitivity functions from laser flash and benactazine (5,6) experiments replotted to reflect loss in contrast sensitivity over the spatial frequency spectrum relative to baseline contrast sensitivity.

Inverse transformed images based on the data from each of these experiments are shown in Figure 2a,b,c,d. The upper left panel is the normal digitized image(2a). In the upper right panel the image has been transformed according to changes induced by acute laser exposure measured at 2 minutes post-exposure. The lower left and right hand panels (c and d) correspond to the benactazine and atropine measured contrast sensitivity changes.

Because the effect of acute laser exposure is nearly uniform over the measured spatial frequency region, the image (b) appears to be reduced in overall contrast, unlike that of the same image transformed for the benactazine deficit (c). In this case, a definite blurring of the image is perceived, as fine spatial frequency loss is relatively greater than losses at the lower spatial frequencies. Minimal change in the perceived image qualities is observed for the atropine (d) condition.




FIGURE 2. Computer generated images.

DISCUSSION

In this paper, we have demonstrated that contrast sensitivity data can be utilized to create degraded images that correspond to the measured changes obtained in spatial vision experiments. With this procedure, complex images can be modified to reflect the visual systems measured response to various noxious environmental conditions. More importantly, with such procedures, any image that can be digitized, can be modified and used in specific training scenarios. This method extends our ability to characterize complex stimuli by allowing both normal high contrast as well as degraded images to be represented along the spatial frequency dimension. The knowledge that certain exposure conditions might blur vision while others might simply alter the overall contrast of a complex scene is information that could only safely be provided with the aid of the present technique for characterizing the effects of noxious environmental conditions on spatial vision.

Finally, for the development of effective combat training, incorporation of the degradation technique provided here offers a viable and safer alternative to training procedures requiring adverse environmental conditions capable of producing moderate transient physiological effects.

REFERENCES

1. Ginsburg AP. On a filter approach to understanding the perception of visual form. In: Recognition of Pattern and Form. Ed. by Duane G. Albrecht, Springer-Verlag, New York, 1982;175-192.
2. Ginsburg AP. Proposed new vision standards for the 1980's and beyond: Contrast Sensitivity. In: AGARD Conference Proceedings No. 310 Aerospace Medical Panel Specialist Meeting, Toronto, Canada 1981.
3. Johnson J. Analysis of image forming systems. In: Proceedings of the Image Intensifier Symposium, U.S. Army Engineer Research and Development Laboratory 1958; 6-7 (Oct):249-273.
4. Monroe D., Zwick H. Introduction to digital image processing. Presidio of San Francisco, Ca: Letterman Army Institute of Research, 1984; Laboratory Note No 85-53.
5. Zwick H, Bloom KR, Lund DJ, Beatrice ES. Laser ocular flash effects. In: Combat Ocular Problems, Suppl, Ed. Edwin S. Beatrice, Presidio of San Francisco, California: Letterman Army Institute of Research, April 1982;45-57.
6. Brown B, Haegerstrom-Portnoy G, Adams AJ, Jones RT, Jampolsky A. Effects of anticholinergic drug, benactyzine hydrochloride, on vision and vision performance. Aviation, Space and Environmental Medicine, 1982;53:759-765.
7. Brown B, Baker R, Adams A, Haegerstrom-Portnoy G, Jones R, Jampolsky A. Investigations of visual performance after administration of cholinergic blocking agents. II: Atropine. U.S. Army Medical Research and Development Command, 1982; Contract No. DAMD 17-80-C-0066.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Laboratory Note No. 86-57	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) COMPUTER VISUAL SIMULATION OF CONTRAST SENSITIVITY DEFICITS INDUCED BY LASER AND CHEMICAL ANTIDOTE EXPOSURE	5. TYPE OF REPORT & PERIOD COVERED	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) Harry Zwick, PhD; Dan Monroe, CPT, MSC; Larry Sherman, BS	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Letterman Army Institute of Research Division of Ocular Hazards Presidio of San Francisco, CA 94129-6800	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS WU 245, Physiologic Basis of Laser Effects, 3M161102BS10 Task: CE	
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research & Development Command Fort Detrick Frederick, MD 21701	12. REPORT DATE	
	13. NUMBER OF PAGES	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report)	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) This report has been approved for sale; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Computer image analysis; Contrast sensitivity; Image simulation; Degraded images; Spatial vision; Noxious environments; Blurred vision; Adverse conditions; Benactazine; Atropine.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Training in some complex combat-related tasks may produce a degree of transient visual impairment which may simulate what could be expected in combat. This paper presents a method for simulating visual impairment produced by potential combat conditions. The use of a computer to both digitize and store as well as produce the simulated image has provided an ideal tool for research. The degree of realism provided by such simulation offers suggestions for development of more realistic training techniques. Key words:		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)